



Health Officer Report Infectious Disease Update

03.08.24

TRACIE NEWMAN, MD, MPH, FAAP



Pertussis Outbreak

- Since October, 39 cases in Cass County (1 border case) and increasing; since Feb 1st, 19 cases
- Cass County averages 5-8 pertussis cases/year
- Ensure you are up to date on pertussis vaccine
- Prophylactic antibiotics
- Suspected and confirmed pertussis cases should exclude form work/school/childcare for 5 days after start
 of meds or 21 days after onset of cough
- See detailed recommendations in our recent FCPH News Release: <u>The City of Fargo News Detail</u> (fargond.gov)

Hepatitis A Case in Grand Forks, ND

- Food worker at Chick-fil-A (UND Memorial Union) likely exposed from travel out of state
- People potentially exposed recommended to receive hep A vaccine / be seen if experiencing symptoms
- NDHHS Health Alert Network Health Advisory sent to providers
- GFPH hosted free hep A vaccine clinic
- To date, no additional cases reported



Hepatitis A

Vaccine-preventable liver infection

Very infectious, caused by hepatitis A virus

Transmission: eating contaminated food/water, person-to-person contact, other high-risk exposures

15-50 days for symptoms to develop

Symptoms can last up to 2 months and include:

- Fatigue
- Nausea
- Stomach pain
- Jaundice

Best prevention = hep A vaccine ASAP (people with 2 doses of hep A vaccine or who have had disease, do not need to vaccinate)

Measles Outbreaks Increasing in U.S.



To date, in 2024, 41 cases in 16 jurisdictions (including MN)



Concerning outbreak in Florida elementary school (at least 8 cases)


Florida State Health Department not upholding standard of practice public health guidelines – allowing unvaccinated exposed children to not isolate for 21 days and not recommending those without immunity receive MMR vaccine

North Dakota Immunization Rates

Choose grade to display Statewide immunization and exemption rates by grade.


*Exemption rates are reported as overall exemption rates for kindergarten and MMR exemption rates for seventh and eleventh.








Coverage Rates for all Required Vaccines




Coverage Rates for all Required Vaccines



State Level Exemption Rates*

● Personal Belief Exemption ● Medical Exemption ● Religious Exemption




State Level Exemption Rates*


● Personal Belief Exemption ● Medical Exemption ● Religious Exemption



Source: NDHHS Disease Control Section





Percent of North Dakota children 19 - 35 months of age by the last day of the quarter who are up-to-date with the selected vaccine by the end of the quarter*



Season: 2023-24 and 5 previous seasons ▼

Percentage of Outpatient Visits for Respiratory Illness Reported by The U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet), Weekly National Summary, 2023-24 Season and Selected Previous Seasons





COVID-19 Update for the United States

Early Indicators



% Test Positivity

8.1%

(February 11 to February 17, 2024)

Trend in % Test Positivity

-1.3% in most recent week

Dec 30, 2023

Feb 17, 2024

Emergency Department Visits

% Diagnosed as COVID-19

1.7%

(February 11 to February 17, 2024)

Trend in % Emergency Department Visits

-12.4% in most recent week

Dec 30, 2023

Feb 17, 2024

These early indicators represent a portion of national COVID-19 tests and emergency department visits. <u>Wastewater</u> information also provides early indicators of spread.

Severity Indicators

Hospitalizations

Hospital Admissions

18,977

(February 11 to February 17, 2024)

Trend in Hospital Admissions

-5.7% in most recent week

Dec 30, 2023

Feb 17, 2024

Total Hospitalizations

6,834,006

Deaths >

% of All Deaths in U.S. Due to COVID-19

2.4%

(February 11 to February 17, 2024)

Trend in % COVID-19 Deaths

-11.1% in most recent week

Dec 30, 2023 Feb 17, 2024

Total Deaths


1,180,042

Nationwide RSV Activity Decreasing

RSV Levels decreasing in many regions

RSV hospitalizations decreased for infants and the elderly

RSV deaths remain stable





2023-2024 Influenza Weekly Report

North Dakota Department of Health and Human Services

Week 9

Ending Saturday, March 02, 2024 Last updated 03/05/2024

Influenza B now makes up the majority of new lab-diagnosed influenza cases in ND. Several metrics of flu activity increase from the previous week, including aggregate laboratory positivity and the percentage of outpatient visits for flu-like illness. Nationwide, seasonal flu activity remains elevated with increases in certain areas of the country.

Beginning January 1st, 2024, NDHHS updated its reportable conditions list and influenza is now reportable via electronic laboratory report (ELR) only. More information detailing this change can be found on our NDHHS website <a href="https://example.com/here-new-nd-new

Everyday preventative actions, including frequent handwashing and covering coughs/sneezes, are easy and effective methods at reducing the spread of influenza and other respiratory diseases this fall. For more information regarding these and more, visit ndflu.com.

Subtype	Cases in Previous Week	Total for Season
Influenza A	0	6,283
Influenza A 2009 H1N1	0	283
Influenza A H3N2	0	82
Influenza B	0	2,198
Influenza B Victoria	0	76
Total	0	8,922

	Last Week	Season Total
New Influenza Cases:	0	8,922
Outpatient Visits for Influenza-like Illness:	0.00%	4.70%
Laboratory Specimens Positive for Influenza:	0.00%	15.28%
Percentage of Students Absent from School:	0.25%	2.06%
New Hospitalizations due to Influenza:	0	339
New Deaths due to Influenza:	0	29





Figure 6. Monthly Nirsevimab Receipt and Intent Among Females Aged 18-49 Years Who Have an Infant<8 Months, Are Currently Pregnant, or Are Trying to Get Pregnant, United States*,†

Data Source: National Immunization Survey–Adult COVID Module




National Vaccine Coverage for COVID-19, Flu, RSV Low. *Good News:

U.S. Nirsevimab stats as of 1/24:

- 40.5% infants < 8 months received
- 21.7% definitely plan to receive
- Pregnant women: 38.2% definitely plan for infant to receive
- Women trying to get pregnant: 44.4% definitely plan to get nirsevimab for infant

ACIP says adults 65+ should get 2nd COVID-19 booster this spring





Questions